ARTHROSCOPIC SLAP REPAIR REHABILITATION PROTOCOL

PHASE I - IMMEDIATE POSTOPERATIVE PHASE (Postoperative Day 1 to Week 6) Goals:

- Protection of anatomic SLAP repair
- Control postoperative pain / inflammation
- Promotion of shoulder musculature contraction
- > Retard muscular atrophy
- > Early controlled motion glenohumeral / scapulothoracic joints

WEEK 0-3

- ➤ Sling 4 weeks or per MD recommendation
- ➤ Sleep in shoulder sling / brace for 4 weeks
- Elbow, wrist, hand ROM, gripping exercise
- PROM / light AAROM glenohumeral joint
 - Flexion to 60° (week1)
 - o Flexion to 70°-75° (week2)
 - o ER to 10°
 - o IR to 30°
- Submaximal isometrics for shoulder musculature (no biceps)
- No active shoulder ER / Extension / Abduction
- Initiate proprioceptive training end of week 1

WEEK 3-4

- ➤ Discontinue sling / immobilizer at 3-4 weeks or per MD recommendation
- Continue PROM / AAROM (progression based on evaluation of patient and subjective patient report)
 - Flexion to 90°
 - o Abduction to 80°
 - o ER to 30°
 - o IR to 40°
- ➤ AAROM: scapular plane elevation and ER to above limits
- Scaption isometrics @ 30 degrees (against gravity)
- Initiate scapular neuromusculature reeducation (rhythmic stabilization)
- Continue shoulder isometrics (no bicep contractions)
- ➤ Initiate shoulder isotonic program at 0° of shoulder abduction ER / IR
- > Cryotherapy with progression to thermal (heat) at week 3

WEEK 4-6

- Gradually improve ROM of shoulder
 - Flexion to 145°



- o ER at 45° abduction: 45°-50°
- o ER at 45° abduction: 55°-60°
- $\circ~$ At 6 weeks begin slow progression toward ER at 90° abduction progress to $30^{\circ}\text{-}40^{\circ}$ ER
- ➤ May initiate stretching exercises
- Continue isotonic tubing ER/IR (arm at side)
- Initiate Active Shoulder Abduction (without resistance)
- Initiate "Full Can" Exercise (Weight of Arm)
- ➤ Initiate Prone Rowing, Prone Horizontal Abduction
- ➤ NO Biceps Strengthening
- ➤ Bodyblade: scaption @ 0°, ER/IR * not aggressive*
- Dynamic weightbearing on ball (bilateral)

Phase II - Intermediate Phase: Moderate Protection Phase (Week 6 - 14) Goals:

- ➤ Gradually restore full ROM (week 10-12)
- Protect the integrity of the surgical repair
- Restore muscular strength and balance

WEEK 6-8

- > Gradually progress to full ROM:
 - o Flexion to 180°
 - o ER at 90° abduction: 90°-95°
 - o IR at 90° abduction: 70°-75°
- Continue to progress isotonic strengthening program
- Continue PNF strengthening
- Initiate Throwers Ten Program
- > ER/IR with resistive tubing
- > ER/IR standing with weight
- > Resisted scaption

WEEK 8-10

- ➤ PROM: to tolerance including cocking (ER @ 90° ABD)
- Bodyblade horizontal abd/add
- Supine cocking with weight (eccentric emphasis)
- Supine PNF D2 diagonal with weight (eccentric emphasis)
- Dynamic weightbearing on ball (unilateral)

WEEK 10+

- May initiate slightly more aggressive strengthening
- Progress ER to Throwers Motion
 - o ER at 90° abduction: 110°-115° in throwers (Week 10-12)
- Progress isotonic strengthening exercises to include manual resistance
- Continue all stretching exercises **Progress ROM to functional demands (i.e. overhead athlete)



- Continue all strengthening exercises
- Rebounder cocking and backhand toss
- > Push ups
- Increase speed of training
- Increase emphasis on eccentric control of cocking

Phase III - Minimal Protection Phase (Week 14 - 20)

Goals:

- Establish and maintain full ROM (throwers motion)
- ➤ IMPROVE muscular strength
- > INITIATE power and endurance. Gradually initiate functional activities

Criteria to enter Phase III:

- > Full non-painful ROM
- Satisfactory stability
- Muscular strength (good grade or better)
- ➤ No pain or tenderness

Week 14 - 16

- Continue all stretching exercises (capsular stretches)
- ➤ Maintain Throwers Motion (emphasis on shoulder ER)
- Continue strengthening exercises:
 - o Throwers Ten Program with progression to Advanced Throwers Program
 - o PNF Manual Resistance
 - o Endurance training (stabilization/perturbation at end range 90/90 position)
 - o Initiate light plyometric program
 - o Restricted sport activities (light swimming, half golf swings)

Week 16 - 20

- Continue all exercise listed above
- Continue all stretching
- Continue Throwers Ten Program / Advanced Throwers 10 program
- Continue Plyometric Program (90/90 positioning plyos)
- Initiate interval sport program (throwing)
- **See interval Throwing Program

Phase IV - Advanced Strengthening Phase (Week 20 - 26)

Coalc

- Enhanced muscular strength, power and endurance
- Progress functional activities
- Maintain shoulder mobility

Criteria to enter Phase IV:

Full non-painful ROM



- > Satisfactory static stability
- ➤ Muscular strength 75-80% of contralateral side
- ➤ No pain or tenderness

Week 20 - 26

- Continue flexibility exercises
- Continue isotonic strengthening program
- > PNF manual resistance patterns
- > Plyometric strengthening
- Progress interval sport programs

Phase V – Return to Activity Phase (Month 6 to 9) Goals:

- Gradual return to sport activities
- > Maintain strength, mobility and stability

Criteria to enter Phase V:

- > Full functional ROM
- Muscular performance isokinetic (fulfills criteria)
- > Satisfactory shoulder stability upon examination
- > No pain or tenderness

Exercises:

- Gradually progress sport activities to unrestrictive participation
- Continue stretching and strengthening program